



The Sparse Cardinal Sine Decomposition (SCSD) and its application to the simulation of suspensions

François Alouges, Matthieu Aussal, Aline Lefebvre-Lepot, Franck Pigeonneau, Antoine Sellier

► To cite this version:

François Alouges, Matthieu Aussal, Aline Lefebvre-Lepot, Franck Pigeonneau, Antoine Sellier. The Sparse Cardinal Sine Decomposition (SCSD) and its application to the simulation of suspensions. International Association for Boundary Element Methods (IABEM), Jun 2018, Paris, France. hal-01978959

HAL Id: hal-01978959

<https://hal-mines-paristech.archives-ouvertes.fr/hal-01978959>

Submitted on 12 Jan 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**The Sparse Cardinal Sine Decomposition (SCSD)
and its application to the simulation of suspensions.**

Francois Alouges¹, Matthieu Aussal¹, Aline Lefebvre-Lepot^{2,*}, Franck Pigeonneau³,
Antoine Sellier⁴

¹CMAP, Ecole Polytechnique, Palaiseau, France

²CNRS, CMAP, Ecole Polytechnique, Palaiseau, France

³CEMEF, MINES ParisTech, Sophia Antipolis, France

⁴LadHyX, Ecole Polytechnique, Palaiseau, France

*Email: aline.lefebvre@polytechnique.edu

Keywords: Boundary-integral equations, SCSD, Stokes equations, Suspensions.

For many applications (settling, transport...), it is necessary to compute the flow of either dilute or concentrated unbounded suspensions made of solid particles immersed in a Newtonian liquid. For small enough particles, the flow Reynolds number vanishes and the task fortunately reduces to the treatment of the linear Stokes equations [1]. This can be efficiently achieved by solving boundary-integral equations [2].

However, as usual in such formulations, the discretization of the problem leads to dense and non-symmetric linear systems whose size grows as the square of the number of particles. Acceleration techniques are therefore usually employed. Most of them are based on the compression of the underlying matrix in order to obtain efficient matrix vector products (see e.g. Fast Multipole Method, \mathcal{H} -matrices, etc.). In this direction, the new Sparse Cardinal Sine Decomposition method (SCSD) was recently developed for the scalar kernel encountered in acoustics [3]. The main idea consists in expanding the kernel in the Fourier space as a finite sum of Cardinal Sine functions. The method has been further extended to the vectorial Stokes kernel in [4] where it has actually been implemented and tested for a single solid particle.

After presenting the SCSD solver for the Stokes kernel, this work investigates its ability to efficiently cope with N -particle clusters immersed in a Newtonian liquid. Cases of large N will be investigated. Both distant or close (packed) particles will be considered.

Acknowledgements

The authors thank Saint-Gobain Research for partially supporting this work.

References

- [1] J. Happel, H. Brenner, *Low Reynolds number hydrodynamics: with special applications to particulate media*, Springer Science & Business Media, 2012.
- [2] C. Pozrikidis, *Boundary Integral and Singularity Methods for Linearized Viscous Flow*, Cambridge University Press: Cambridge, 1992.
- [3] F. Alouges, M. Aussal, *The sparse cardinal sine decomposition and its application for fast numerical convolution*, Numerical Algorithms, **70**(2), pp. 427–448, 2015.
- [4] F. Alouges, M. Aussal, A. Lefebvre-Lepot, F. Pigeonneau, A. Sellier, *Application of the sparse cardinal sine decomposition applied to 3D Stokes flows*, International Journal of Comp. Meth. and Exp. Meas., **vol 5**(3), pp. 387–394, 2017.